SEP 2 5 2006

Reply under 37 CFR 1.116
Expedited Procedure
Technology Center 1624
Attorney Docket No. CV06038US01

AMENDMENTS TO THE CLAIMS

1. (Currently Amended) A compound represented by the structural formula (I):

$$Q^{1} \qquad Q^{2} \qquad Q^{3} \qquad Q^{3} \qquad Q^{3} \qquad Q^{4} \qquad Q^{5} \qquad Q^{5} \qquad Q^{4} \qquad Q^{4$$

or pharmaceutically acceptable isomers, salts, solvates or esters of the compound of Formula (I), wherein in Formula (I) above:

X, Y and Z can be the same or different and each is independently selected from the group consisting of -CH₂-, -CH(alkyl)- and -C(alkyl)₂-;

 Q^1 and Q^2 can be the same or different and each is independently selected from the group consisting of H, $\underline{-G_1}$ - $(C_{01}$ - C_{30} alkylene)-G, $-OR^6$, $-OC(O)R^6$, $-OC(O)OR^9$,

 $-OC(O)NR^6R^7$ and -L-M;

Q³ is 1 to 5 substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, -G, -(C₁-C₃₀ alkylene)-G, -OR⁶, -(C₁-C₁₀ alkylene)-OR⁶, -C(O)R⁶,

 $-(C_1-C_{10} \text{ alkylene})-C(O)R^6$, $-C(O)OR^6$, $-(C_1-C_{10} \text{ alkylene})-C(O)OR^6$, $-OC(O)R^6$,

-(C₁-C₁₀ alkylene)-OC(O)R⁶, -OC(O)OR⁹, -(C₁-C₁₀ alkylene)-OC(O)OR⁹, -CH=CH-C(O)R⁶,

 $-CH=CH-C(O)OR^6$, $-C=C-C(O)OR^6$, $-C=C-C(O)R^6$, $-O-(C_1-C_{10} \text{ alkylene})-OR^6$,

 $-O-(C_1-C_{10} \text{ alkylene})-C(O)R^6$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)OR^6$, -CN,

 $-O-(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7$, $-O-C(O)NR^6NR^7C(O)OR^6$,

-O-(C_1 - C_{10} alkylene)-C(O)NR⁶NR⁷C(O)OR⁶, -O-(C_1 - C_{10} alkylene)-C(O)(aryl)-N₃,

 $-OC(O)-(C_1-C_{10} \text{ alkylene})-C(O)OR^6$, $-C(O)NR^6R^7$, $-(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7$,

 $-OC(O)NR^6R^7$, $-(C_1-C_{10} \text{ alkylene})-OC(O)NR^6R^7$, $-NO_2$, $-NR^6R^7$. $-(C_1-C_{10} \text{ alkylene})-NR^6R^7$,

-O-(C₂-C₁₀ alkylene)-NR⁶R⁷, -NR⁶C(O)R⁷, -NR⁶C(O)OR⁹,
-NR⁶C(O)NR⁷R⁸, -NR⁶S(O)₀₋₂R⁹, -N(S(O)₀₋₂R⁹)₂, -CHNOR⁶, -C(O)NR⁶R⁷,
-C(O)NR⁶NR⁶R⁷, -S(O)₀₋₂NR⁶R⁷, -S(O)₀₋₂R⁹, -O-C(O)-(C₁-C₁₀ alkylene)-C(O)NR⁶R⁷,
-OC(O)-(C₁-C₁₀ alkylene)-NR⁶C(O)O-(alkylaryl), -P(O)(OR¹⁰)₂,
-(C₁-C₁₀ alkylene)-OSi(alkyl)₃, -CF₃, -OCF₃, halo, alkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyarylalkoxy, alkoxyarylalkoxy, alkoxyarylalkoxy, alkoxyarylalkoxy, alkoxyarylalkoxy, arylalkoxy, arylalkoxy, arylalkoxy, arylalkoxy, arylalkoxy, arylalkoxy, arylalkoxy, arylalkoxy, arylalkoxy, heteroarylalkyl, heteroarylalkoxy, dioxolanyl, heterocyclylalkyl, heterocyclylcarbonyl, heterocyclylcarbonylalkoxy and -L-M;

Q4 is 1 to 5 substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, -G, -(C_1 - C_{30} alkylene)-G, -OR⁶, -(C_1 - C_{10} alkylene)-OR⁶, -C(O)R⁶, - $(C_1-C_{10} \text{ alkylene})-C(O)R^6$, - $C(O)OR^6$, - $(C_1-C_{10} \text{ alkylene})-C(O)OR^6$, - $OC(O)R^6$, -(C₁-C₁₀ alkylene)-OC(O)R⁶, -OC(O)OR⁹, -(C₁-C₁₀ alkylene)-OC(O)OR⁹, -CH=CH-C(O)R⁶, $-CH=CH-C(O)OR^6$, $-C=C-C(O)OR^6$, $-C=C-C(O)R^6$, $-O-(C_1-C_{10} \text{ alkylene})-OR^6$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)R^6$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)OR^6$, -CN, $-O-(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7$, $-O-C(O)NR^6NR^7C(O)OR^6$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)NR^6NR^7C(O)OR^6$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)(aryl)-N_3$, $-OC(O)-(C_1-C_{10} \text{ alkylene})-C(O)OR^6$, $-C(O)NR^6R^7$, $-(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7$, $-OC(O)NR^6R^7$, $-(C_1-C_{10} \text{ alkylene})-OC(O)NR^6R^7$, $-NO_2$, $-NR^6R^7$, $-(C_1-C_{10} \text{ alkylene})-NR^6R^7$, $-O-(C_2-C_{10} \text{ alkylene})-NR^6R^7, -NR^6C(O)R^7, -NR^6C(O)OR^9, -NR^6C(O)NR^7R^8, -NR^6S(O)_{0.2}R^9, -NR^6C(O)NR^7R^8, -NR^6S(O)_{0.2}R^9, -NR^6C(O)NR^7R^8, -NR^6S(O)_{0.2}R^9, -NR^6C(O)NR^7R^8, -NR^6S(O)_{0.2}R^9, -NR^6C(O)NR^7R^8, -NR^6S(O)_{0.2}R^9, -NR^6C(O)NR^7R^8, -NR^6S(O)_{0.2}R^9, -NR^6C(O)_{0.2}R^9, -NR^6C(O)$ $-N(S(O)_{0.2}R^9)_2$, $-CHNOR^6$, $-C(O)NR^6R^7$, $-C(O)NR^6NR^6R^7$, $-S(O)_{0.2}NR^6R^7$, $-S(O)_{0.3}R^9$. $-O-C(O)-(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7$, $-OC(O)-(C_1-C_{10} \text{ alkylene})-NR^6C(O)O-(\text{alkylaryl})$, $-P(O)(OR^{10})_2$, $-(C_1-C_{10} \text{ alkylene})-OSi(alkyl)_3$, $-CF_3$, $-OCF_3$, halo, alkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxycarbonylalkoxy, alkoxyarylalkoxy, alkoxyiminoalkyl, alkyldioyl, allyloxy, aryl, arylalkyl, aryloxy, arylalkoxy, aroyl, aroyloxy, aroylaroyloxy, arylalkoxycarbonyl,

benzoylbenzoyloxy, heteroaryla, heteroarylalkyl, heteroarylalkoxy, dioxolanyl, heterocyclylalkyl, heterocyclylcarbonyl, heterocyclylcarbonylalkoxy and -L-M;

 Q^5 is 1 to 5 substituents independently selected from the group consisting of alkyl, alkynyl, -G, -(C₁-C₃₀ alkylene)-G, -OR⁶, -(C₁-C₁₀ alkylene)-OR⁶, -C(O)R⁶,

- $(C_1-C_{10} \text{ alkylene})$ - $C(O)R^6$, - $C(O)OR^6$, - $(C_1-C_{10} \text{ alkylene})$ - $C(O)OR^6$, - $OC(O)R^6$,

- $(C_1-C_{10} \text{ alkylene})$ - $OC(O)R^6$, - $OC(O)OR^9$, - $(C_1-C_{10} \text{ alkylene})$ - $OC(O)OR^9$, - $CH=CH-C(O)R^6$,

 $-CH=CH-C(O)OR^6$, $-C==C-C(O)OR^6$, $-C==C-C(O)R^6$, $-O-(C_1-C_{10}$ alkylene)-OR⁶,

 $-O-(C_1-C_{10} \text{ alkylene})-C(O)R^6$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)OR^6$, -CN,

-O-(C_1 - C_{10} alkylene)-C(O)NR⁶R⁷, -O-C(O)NR⁶NR⁷C(O)OR⁶,

 $-O-(C_1-C_{10} \text{ alkylene})-C(O)NR^6NR^7C(O)OR^6$, $-O-(C_1-C_{10} \text{ alkylene})-C(O)(aryl)-N_3$,

 $-OC(O)-(C_1-C_{10} \text{ alkylene})-C(O)OR^6, -C(O)NR^6R^7, -(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7, -(C_1-C_1) \text{ alkylene}$

 $-OC(O)NR^6R^7$, $-(C_1-C_{10} \text{ alkylene})-OC(O)NR^6R^7$, $-NO_2$, $-NR^6R^7$, $-(C_1-C_{10} \text{ alkylene})-NR^6R^7$,

-O-(C2-C10 alkylene)-NR⁶R⁷, -NR⁶C(O)R⁷, -NR⁶C(O)OR⁹,

 $-NR^6C(O)NR^7R^8$, $-NR^6S(O)_{0.2}R^9$, $-N(S(O)_{0.2}R^9)_{2}$, $-CHNOR^6$, $-C(O)NR^6R^7$,

 $-C(O)NR^6NR^6R^7$, $-S(O)_{0.2}NR^6R^7$, $-S(O)_{0.2}R^9$, $-O-C(O)-(C_1-C_{10} \text{ alkylene})-C(O)NR^6R^7$,

 $-OC(O)-(C_1-C_{10} \text{ alkylene})-NR^6C(O)O-(\text{alkylaryl}), -P(O)(OR^{10})_2,$

-(C₁-C₁₀ alkylene)-OSi(alkyl)₃, -CF₃, -OCF₃, halo, alkoxyalkoxy, alkoxyalkoxy, alkoxyarbonylalkoxy, alkoxyarylalkoxy, alkoxyiminoalkyl, alkyldioyl, allyloxy, arylalkyl, aryloxy, arylalkoxy, aroylaroyloxy, arylalkoxycarbonyl, benzoylbenzoyloxy, heteroarylalkyl, heteroarylalkoxy, dioxolanyl, heterocyclylalkyl, heterocyclylcarbonyl, heterocyclylcarbonyl, heterocyclylcarbonylalkoxy and -L-M;

wherein optionally one or more carbon atoms of the $-(C_1-C_{30} \text{ alkylene})$ - radical of Q^1 , Q^2 , Q^3 , Q^4 and Q^5 is independently replaced by -O-, -C(O)-, -CH=CH-, -C=C-, -N(alkyl)-, -N(alkyl)-, or -NH-;

G is selected from the group consisting of a sugar residue, disugar residue, trisugar residue, tetrasugar residue, sugar acid, amino sugar, amino acid residue, oligopeptide residue

comprising 2 to 9 amino acids, trialkylammoniumalkyl saltradical and -S(O)₂-OH, wherein optionally the sugar residue, disugar residue, trisugar residue, tetrasugar residue, sugar acid, amino sugar, amino acid residue or oligopeptide residue of G is substituted with -L-M;

L is selected from the group consisting of

$$\begin{array}{c} \cdot \xi - O - C(O) & \\ \cdot \xi - O - C(O) - (CH_2)_{xx1} - (O)C - \xi \\ \cdot \xi - O - C(O) - (CH_2)_{xx3} - (O)C - \xi \\ \cdot \xi - O - SiMe_2 - (CH_2)_{xx3} - (O)C - \xi \\ \cdot \xi - O - SiMe_2 - (CH_2)_{xx3} - OC(O) - \xi \\ \cdot \xi - O - SiMe_2 - (CH_2)_{xx3} - OC(O) - \xi \\ \cdot \xi - O - SiMe_2 - (CH_2)_{xx3} - OC(O) - \xi \\ \cdot \xi - O - C(O) - (CH_2)_{xx3} - OC(O) - \xi \\ \cdot \xi - O - C(O) - (CH_2)_{xx10} - OC(O) - \xi \\ \cdot \xi - OC(O) - (CH_2)_{xx10} - OC(O) - \xi \\ \cdot \xi - OC(O) - (CH_2)_{xx10} - OC(O) - \xi \\ \cdot \xi - OC(O) - (CH_2)_{xx10} - OC(O) - \xi \\ \cdot \xi - OC(O) - (CH_2)_{xx10} - OC(O) - \xi \\ \cdot \xi - OC(O) - (CH_2)_{xx10} - OC(O) - \xi \\ \cdot \xi - OC(O) - (CH_2)_{xx10} - OC(O) - \xi \\ \cdot \xi - OC(O) - (CH_2)_{xx10} - OC(O) - \xi \\ \cdot \xi - OC(O) - (CH_2)_{xx10} - OC(O) - \xi \\ \cdot \xi - OC(O)$$

wherein Me is methyl;

M is selected from the group of moieties consisting of

$$H_2C$$
 CH_2
 H_3C
 CH_3
 CH_3

$$\begin{array}{c} \text{OH} \\ \\ \text{H}_2\text{O} \\ \text{CH}_2 \\ \\ \text{CH}_3 \\ \text{CH}_3 \\ \text{CH}_4 \\ \\ \text{CH}_5 \\ \\ \text{CH}_5 \\ \\ \text{CH}_5 \\ \\ \text{CH}_5 \\ \\ \text{CH}_6 \\ \\ \text{CH}_6 \\ \\ \text{CH}_6 \\ \\ \text{CH}_7 \\ \\ \text{CH}_8 \\ \\ \text{CH$$

pharmaceutically acceptable salts of the moieties (M1) to (M9) and free acids of the moieties (M1) to (M9);

R² and R³ can be the same or different and each is independently selected from the group consisting of hydrogen, alkyl and aryl;

R⁶, R⁷ and R⁸ can be the same or different and each is independently selected from the group consisting of hydrogen, alkyl, aryl and arylalkyl; and

each R⁹ is independently alkyl, aryl or arylalkyl.

each R¹⁰ is independently H or alkyl;

q is 0 or 1;

r is 0 or 1;

m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3, 4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5;

x1 is 1 to 10;

x2 is I to 10;

x3 is 1 to 10;

x4 is 1 to 10;

x5 is 1 to 10;

x6 is 1 to 10; and

x7 is 1 to 10;

x8 is 1 to 10;

x9 is 1 to 10; x10 is 1 to 10; and x11 is 1 to 10;

with the proviso that at least one of Q¹, Q², Q³, Q⁴ and Q⁵ is –L-M or the sugar residue, disugar residue, trisugar residue, tetrasugar residue, sugar acid, amino sugar, amino acid residue or oligopeptide residue of G is substituted with –L-M, and wherein each of the above alkyl, alkenyl, alkynyl, alkylene, alkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyarylalkoxy, alkoxyarylalkoxy, alkoxyiminoalkyl, alkyldioyl, allyloxy, aryl, arylalkyl, aryloxy, arylalkoxy, aroylaroyloxy, arylalkoxycarbonyl, benzoylbenzoyloxy, heteroarylalkyl, heteroarylalkoxy, dioxolanyl, heterocyclylalkyl, heterocyclylcarbonyl, or heterocyclylcarbonylalkoxy groups, when present, is independently substituted or unsubstituted.

- 2. (Original) The compound according to claim 1, wherein m, n and r are each zero, q is 1, p is 2, and Z is -CH₂-.
- 3. (Original) The compound according to claim 1, wherein m, n and r are each zero, q is 1, p is 2, and Z is -CH₂-, Q¹ is -OR⁶, wherein R⁶ is hydrogen and Q⁵ is fluorine.
- 4. (Previously Presented) The compound according to claim 1, wherein R² and R³ are each hydrogen.
- 5. (Original) The compound according to claim 1, wherein Q¹ and Q² are each independently selected from the group consisting of -OR⁶; -O(CO)R⁶, -O(CO)OR⁹ and -O(CO)NR⁶R⁷.
 - 6. (Original) The compound according to claim 1, wherein Q⁴ is halo or -OR⁶.

- (Original) The compound according to claim 1, wherein Q¹ is -OR⁶ wherein R⁶ is

 H.
- 8. (Original) The compound according to claim 1, wherein Q^1 , Q^2 , Q^3 , Q^4 or Q^5 is -L-M.
- 9. (Previously Presented) The compound according to claim 1, wherein Q^1 , Q^2 , Q^3 , Q^4 or Q^5 is -G or $-(C_1-C_{30}$ alkylene)-G.
- 10. (Withdrawn) The compound according to claim 1, wherein G is selected from the group consisting of:

$$R^{4b}Q$$
 R $R^{4b}Q$ R $R^{4b}Q$ R $R^{4b}Q$ $R^{4a}Q$ $R^{4a}Q$

wherein R, R^a and R^b can be the same or different and each is independently selected from the group consisting of H, -OH, halo, -NH₂, azido, alkoxyalkoxy or -W-R³⁰;

W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(\mathbb{R}^{31})-, -NH-C(O)-N(\mathbb{R}^{31})- and -O-C(S)-N(\mathbb{R}^{31})-;

R^{2a} and R^{6a} can be the same or different and each is independently selected from the group consisting of H, alkyl, acetyl, aryl and arylalkyl;

R^{3a}, R^{4a}, R^{5a}, R^{7a}, R^{3b} and R^{4b} can be the same or different and each is independently selected from the group consisting of H, alkyl, acetyl, arylalkyl, -C(O)alkyl and -C(O)aryl;

R³⁰ is independently selected from the group consisting of R³²-substituted T, R³²-substituted-T-alkyl, R³²-substituted-alkenyl, R³²-substituted-cycloalkyl and R³²-substituted-cycloalkylalkyl;

R³¹ is independently selected from the group consisting of H and alkyl;

T is independently selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;

R³² is 1 to 3 substituents which are each independently selected from the group consisting of H, halo, alkyl, -OH, phenoxy, -CF₃, -NO₂, alkoxy, methylenedioxy, oxo, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, -N(CH₃)₂, -C(O)-NHalkyl, -C(O)-N(alkyl)₂, -C(O)-alkyl, -C(O)-alkoxy and pyrrolidinylcarbonyl; or R³² is a covalent bond and R³¹, the nitrogen to which it is attached and R³² form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolinyl or morpholinyl group, or a alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyl or morpholinyl group.

11. (Withdrawn) The compound according to claim 10, wherein G is selected from:

wherein Ac is acetyl and Ph is phenyl.

- 12. (Previously Presented) The compound according to claim 1, wherein optionally one or more carbon atoms of the $-(C_1-C_{30}$ alkylene)- radical of Q^1 , Q^2 , Q^3 , Q^4 and Q^5 is independently replaced by -O -.
- 13. (Original) The compound according to claim 1, wherein L is $-\frac{1}{2}$ O-C(O)-(CH₂)_{x1} (O)C $-\frac{1}{2}$
- 14. (Original) The compound according to claim 1, wherein L is ξ —O—(CH₂) \overline{x} 3 (O)C— ξ

15. (Original) The compound according to claim 1, wherein M is

(M1) or pharmaceutically acceptable salts thereof.

16 (Original) The compound according to claim I, wherein M is

(M2) or pharmaceutically acceptable salts thereof.

17. (Original) The compound according to claim 1, wherein M is

(M3) or pharmaceutically acceptable salts thereof.

18. (Original) The compound according to claim 1, wherein M is

19. (Original) The compound according to claim 1, wherein M is

20. (Original) The compound according to claim 1, which is selected from the group consisting of

- 21. (Currently Amended) A pharmaceutical composition for the treatment of atherosclerosis, hypercholesterolemia, sitosterolemia, diabetes mellitus, obesity, stroke, lowering a concentration of cholesterol, phytosterol or 5α-stanol in plasma of a mammal, treating domyclination or treating Alzhoimer's disease and/or regulating levels of amyloid β peptides in a subject comprising a therapeutically effective amount of a compound of claim 1 in a pharmaceutically acceptable carrier.
- 22. (Original) A pharmaceutical composition comprising a cholesterol-lowering effective amount of a compound of claim 1 in a pharmaceutically acceptable carrier.

- 23. (Currently Amended) A method of treating atherosclerosis, hypercholesterolemia, sitosterolemia, diabetes mellitus, obesity, stroke, lowering a concentration of cholesterol, phytosterol or 5α-stanol in plasma of a mammal, treating demyolination or treating Alzheimer's disease or regulating a level of an amyloid β peptide in a subject comprising the step of administering to a subject in need of such treatment an effective amount of a compound of claim 1.
- 24. (Original) A method of lowering cholesterol level in plasma of a mammal in need of such treatment comprising administering a pharmaceutically effective amount of the compound of claim 1.